Product Name: APVMA Approval No: Y-TEX WARRIOR INSECTICIDAL CATTLE EAR TAGS 51524/119797



Label Name:	Y-TEX WARRIOR INSECTICIDAL CATTLE EAR TAGS
Signal Headings:	POISON
	KEEP OUT OF REACH OF CHILDREN
	FOR ANIMAL TREATMENT ONLY
	READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	100 g/kg CHLORPYRIFOS 300 g/kg DIAZINON (anticholinesterase compounds)
----------------------------	------------------------------------------------------------------------------

Claims:	One tag/head treatment for up to 12 weeks control of diazinon-susceptible Buffalo Flies* and control of cattle lice on beef and dairy cattle (including lactating dairy cows). *Including pyrethroid-resistant strains.
	Resistance may develop to any chemical.

Net Contents:	20 tags - each 15 g [300 g NET weight] 100 tags - each 15 g [1.5 kg NET weight] 500 tags - each 15 g [7.5 kg NET weight] 1000 tags - each 15 g [15.0 kg NET weight] 1300 tags - each 15 g [19.5 kg NET weight]
	It is illegal to sell individual tags separately.

	Directions for Use:					
--	---------------------	--	--	--	--	--

Restraints:	Remove tags prior to slaughter.

Contraindications:	This product is contraindicated for use in calves less than 3 months of age as ear damage
	may occur.

utions:	
---------	--

Side Effects:	

Dosage and Administration:	<dosage administration="" and="" table=""> This section contains file attachment.</dosage>	
-------------------------------	------------------------------------------------------------------------------------------------	--

General Directions:	Y-Tex Warrior Insecticidal Cattle Ear Tags will give effective control of Buffalo Flies and cattle lice if used regularly on all animals in the group (over 3 months of age).
	The treatment regime with a periodic rotation program will reduce the potential for insecticide resistance to develop in Buffalo Flies.
	Application: Apply tags with the Y-TEX UltraTagger® (Figures 1 to 4).
	Figure 1 <drawing> Seat male button firmly on pin, press down and give a slight twist.</drawing>
	Figure 2 <drawing> Place female tag firmly under clip by depressing lever. Collar on tag must be pointing down. Tag and button are now ready for application.</drawing>
	Figure 3 <drawing> Position male stud between the second and third radial cartilage on the front of the ear. Do not allow shaft of the male button to penetrate any cartilage rib or blood vessels or ear damage may result.</drawing>
	Figure 4 <drawing> Squeeze handles of applicator together. When properly positioned, the tag should hang as shown.</drawing>
	NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Withholding Periods:	MEAT: Zero (0) days MILK: Zero (0) days
----------------------	--------------------------------------------

Trade Advice:	TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days. Before using this product, confirm the current ESI from Landmark Operations Limited on 1800 448 892 or the APVMA website www.apvma.gov.au/residues.
---------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Safety Directions:Poisonous if swallowed. Repeated minor exposure may have a cumulative poisoning effect May irritate the eyes. Avoid contact with eyes and skin. Do not open inner pouch until ready for use. Do not allow children to play with tags. When using the product, wear rubber gloves. After use and before eating and drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves.	Safety Directions:
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------

First Aid Instructions:	If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre (Phone Australia 131 126) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.
-------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

|--|

Additional User Safety:	ADDITIONAL USER SAFETY INFORMATION:
	Additional information is listed in the Safety Data Sheet.

Environmental	ENVIRONMENTAL PROTECTION: Use tags only on cattle. DO NOT contaminate streams,
Statements:	rivers or waterways with the tags or used containers.

specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or spent tags.	Disposal:	
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------	--

Storage:	Store below 30°C (Room temperature). Store in original container in a dry, cool place. DO NOT store in direct sunlight.

Y-Tex Warrior Insecticidal Cattle Ear Tags

Dosage and Administration

Animal and Pest	State	Rate	Critical Comments
Cattle (including lactating dairy cattle) Buffalo Flies (<i>Haematobia irritans</i> <i>exigua</i>)	All States and Territories	One tag per animal (1/head)	For Buffalo Fly control: Apply only when fly numbers exceed acceptable levels. <u>Removal of tags after 12 weeks will ensure</u> that Buffalo Fly populations are not exposed to sublethal doses which may increase the chance of resistance. Remove tags with tag removal knife. The level and duration of control may be reduced under high fly pressure. The whole herd is to be treated concurrently.
Cattle Lice (Bovicola bovis, Linognathus vituli, Haematopinus eurysternus, Solenoptes capillatus)			For Cattle Biting Lice control: Apply tags in late Autumn or early Winter and <u>remove after 3 months</u> .